
ALLERGY and Relief from Allergies with the Help of Iodine

Human allergies occur because the body mistakenly identifies harmless substances as enemies and begins attacking the cells where these harmless substances have entered. Thus, there is a failure in the body's "self vs. non-self" recognition system for external substances.

1. How does the immune system determine "self" or "non-self"?

The immune system works through:

- Receptors on cells created according to DNA code
- These receptors recognize antigens (unique fragments of pathogens — proteins, carbohydrates, etc.)
- Memory is stored in B- and T-cells trained during past encounters — a kind of biological RAM + cache, analogous to working memory

2. Allergy — a consequence of an error in immune training

- Normally: the immune system learns in childhood (and partly in the womb) that pollen, food, animal dander, etc. are harmless
- In allergy: this process malfunctions, and the immune system begins to consider harmless substances as enemies
- This occurs because the training of T-cells and B-cells was incorrect or environmental factors contributed to the failure

3. What is the role of DNA and RNA?

- DNA determines how immune cell receptors are built and includes variability via V(D)J recombination (like assembling random locks for future pathogen "keys")
- RNA transmits instructions for synthesizing proteins, including antibodies and receptors

So:

- DNA — generator of recognition templates
- RNA — executor of those templates
- Memory (of friends and enemies) — stored in trained cells, not in DNA

4. Allergy = recognition failure, not absence of knowledge

The immune system “knows” the external substance is not a pathogen but reacts as if it were dangerous.

It is like a security system deciding that a plush toy is a terrorist simply because of a firmware error.

Normal functioning of T-cells and B-cells indeed depends on levels of T3 (triiodothyronine) and T4 (thyroxine) — the main thyroid hormones.

How this is connected:

- **T-cells** (especially in the thymus) are highly sensitive to thyroid hormones
  ➔ T3 and T4 stimulate maturation and differentiation of T-lymphocytes. With hormone deficiency (e.g., hypothyroidism), the thymus shrinks and T-cell maturation is impaired.
- **B-cells** also depend on thyroid hormones, but indirectly — through overall immune regulation and cytokine levels
  ➔ In hypothyroidism, antibody production by B-cells often decreases

If thyroid hormones are low:

- Immune response worsens
- Proliferation of T- and B-cells slows
- Increased susceptibility to infections and reduced vaccine effectiveness

If hormones are too high (hyperthyroidism):

- The immune system may become hyperactive

- Increased risk of autoimmune diseases (e.g., Graves' disease)

Summary Scheme

T3 / T4 (thyroid hormones)
↓
Thymus
➔ Stimulate development and maturation of T-cells
➔ Increase T-cell proliferation
➔ Enhance activity of T-helper (Th1/Th2) and cytotoxic T-cells

↓
Bone marrow and lymph nodes
➔ Support normal B-cell differentiation
➔ Improve antibody production (IgG, IgA, etc.)

↓
Overall immunity
➔ Normal: strong, balanced immune response
➔ Deficiency (hypothyroidism): weakened immunity, infection risk
➔ Excess (hyperthyroidism): autoimmune risk

Visual Mini-Table

T3/T4 Level	Effect on T- and B-cells	Immune Outcome
Normal	Healthy maturation, activation, balance	Strong immunity
Low	Delayed maturation, reduced activity	Weakened immunity

T3/T4 Level	Effect on T- and B-cells	Immune Outcome
High	Hyperactivation, possible autoimmune disruption	Autoimmune risk

---

Cytokines and Receptors Involved in T3/T4 Action

🔵 Major cytokines stimulated by T3 and T4:

Cytokine	Role in immunity	Effect of T3/T4
IL-2	Activates T-cell growth and division	Increased synthesis at normal levels
IFN-γ	Enhances defense against viruses and tumors	Increased production
IL-4	Promotes B-cell maturation into plasma cells	Slightly stimulated
TNF-α	Inflammatory response	May rise sharply in hyperthyroidism

---

🧬 Key receptors through which T3/T4 act:

Receptor	Location	Function
Thyroid hormone receptor α (TRα)	T-cells, B-cells, macrophages	Activates genes for growth, survival, activation
Thyroid hormone receptor β (TRβ)	Antigen-presenting cells	Modulates immune responses and cytokine production

---

✨ Mechanism of Action

1. T3 enters T- and B-cells

---

2. Binds to nuclear receptors TRα or TRβ
3. Activates specific genes, such as:
    - T-cell growth genes
    - IL-2 and IFN-γ secretion genes
4. Immune defense strengthens

If T3/T4 are low → immune processes slow down
If too high → excessive activation → inflammation or autoimmunity

In short:
T3/T4 → Receptor → Genes → Cytokines → Strong immunity 🚀

T4 vs. T3

- T4 — mainly a precursor (“inactive form”)
- T3 — active form that enters cells and binds receptors

T4 converts to T3 via the enzyme deiodinase (removal of one iodine atom).

Without T4 → T3 conversion, immune cells do not receive activation signals.

📚 Iodine and DNA Protection

Iodine (I_2 or iodide) participates not only in thyroid function but also in DNA protection:

1. Antioxidant properties

- Neutralizes free radicals that damage DNA
- Molecular iodine (I_2) is especially effective

2. Direct DNA protection

- Stabilizes DNA under oxidative stress
- Reduces strand breaks in experimental models

3. Activation of “repair genes”

May activate genes responsible for:

- DNA repair
- Antioxidant defense
- Apoptosis of damaged cells

4. Thyroid function and DNA

Iodine deficiency impairs thyroid hormones, which regulate metabolism, cell division, and repair.

🔥 Additional Important Points

- Molecular iodine (I_2) often shows better DNA-protective effects than iodide alone
- Particularly active in tissues with high cell proliferation:
  ➔ Breast, thyroid, stomach, ovaries

⚡ About Dr. Guy Abraham

He promoted the idea that iodine doses above RDA may:

- Improve overall health
- Help cells combat DNA damage
- Possibly reduce cancer risk

---

iod

telegram «iod.kz»
Instagram «iod.kz»
youtube «йод»
tiktok iod.kz
email: info@iod.kz
сайт www.iod.kz
телеграм: +7-777-0554400

📖 What Iodine Really Does for DNA

✅ Helps protect DNA from new damage
✅ Activates natural repair mechanisms
✅ Reduces mutation rates

❗ BUT:

Severely damaged DNA cannot be “magically” restored by iodine alone.

If damage is too severe:

- Repair enzymes attempt correction
- If impossible → apoptosis (programmed cell death)

🔥 Key Conclusion

Stage	Role of Iodine
Protection	Prevents new DNA damage
Mild damage	Supports repair enzymes
Severe damage	Promotes removal of irreparable cells

🧬 Genome Integrity Support

Iodine helps:

- Reduce mutation burden
- Ensure new cells form without inherited damage
- Accelerate tissue regeneration

Iodine does not fix irreversible defects but creates conditions for healthier cell replacement.

📈 Mechanism of Health Restoration

Iodine (I_2 / I^-)
↓
Activation of DNA repair enzymes (PARP, p53, BRCA1)
↓
Damage assessment
➔ Mild → Repair
➔ Severe → Apoptosis
↓
Healthy cell population

In short:
Iodine → Repair or Apoptosis → Tissue renewal 🌱

Iodine acts like a “genome guardian,” helping the body decide which cells to repair and which to remove.

🧬 5 Key DNA Repair Enzymes Influenced by Iodine

1. **PARP (Poly ADP-Ribose Polymerase)**
   Initiates DNA repair by detecting strand breaks
2. **p53 (Tumor Suppressor Protein)**
   Controls cell cycle, repair, or apoptosis
3. **BRCA1**
   Repairs double-strand breaks and maintains genome stability
4. **DNA-PK**
   Repairs double-strand DNA damage (e.g., radiation damage)
5. **ATM (Ataxia Telangiectasia Mutated)**
   Coordinates cellular response to DNA damage

📌 How Iodine Activates These Enzymes

1. Iodine enters cells
2. Interacts with cellular receptors
3. Activates transcription of repair genes
4. Repair enzymes restore DNA integrity

Final Summary

Iodine plays an important role in supporting the cellular DNA repair system by activating enzymes such as PARP, p53, BRCA1, and others. This helps cells recover faster from damage, reduce mutations, and lower cancer risk.